

# Using Geo-Targeted Health Alerts to Respond to Local STI Surges Among Gay and Bisexual Men

<u>Joe Rich</u>, New Zealand AIDS Foundation Jane Morgan, Waikato District Health Board Sunita Azariah. Auckland District Health Board

# Background


Design

Results


Next Steps

## The concentration of Syphilis among MSM

2.5% of NZ population




83% of Syphilis notified in NZ, 2011\*




<sup>\*</sup> Infectious syphilis notified at sexual health clinics. Source: Psutka, R. et al. Int J STD AIDS 2013; 24: 791-798

#### GBM diagnosed with an STI in last 12 months\*







# Condom use during anal intercourse with casual partners among GBM



Saxton, P. Changes in behaviour over 30 years. 8 May 2015.


## Syphilis cases in Waikato SHC





## Syphilis cases in Auckland SHCs

Infectious syphilis diagnosed at Sexual Health Clinics in Auckland\*



#### <u>Objective</u>

Prompt syphilis screening to reduce undiagnosed infections among gay, bisexual and other men who have sex with men (GBM) in Waikato and Auckland.

# Background

Design

Results

Next Steps

#### Health alert to MSM

Syphilis is on the rise

Information about risks and symptoms

Encouraged to use condoms for anal sex and to get tested

#### <u>Channels</u>


Gay news sites: Gaynz.com, Express, Eikon

Hook-up apps: Grindr, Hornet, Growlr, Scruff, Squirt

Porn site: Xtube









# Background

Design

Results

Next Steps


#### Reach

829,173 advertising impressions


4594 clicks to website


#### Syphilis tests among males in Waikato DHB

and GPs)





## Syphilis cases in Waikato SHC






## Syphilis cases in Waikato SHC







## Syphilis cases in Auckland SHCs

Infectious syphilis
diagnosed at Sexual
Health Clinics in
Auckland\*




## Syphilis cases in Auckland SHCs





#### Syphilis cases in New Zealand SHCs





#### Conclusions

Geo-targeted health alerts are an effective way of reaching GBM.

Data on GBM testing numbers is not available so we can't tell to what extent the increase in diagnoses is a result of increased screening.

Patients reported seeing syphilis health alerts to clinicians.

# Background

Design

Results

Next Steps

#### Improve reporting

DHB reporting on GBM testing numbers.

Consider asking "what prompted your test today" in pretest questionnaire; operational and reporting implications.

SHC data does not include ID and GP cases. Making syphilis notifiable will provide a more complete picture.


#### Collaborate

NZAF has the expertise to reach GBM and prompt STI screening.

Driving HIV testing among high-risk GBM can have secondary impact on increasing syphilis screening.

Consider extending syphilis rapid testing alongside HIV rapid testing in primary care settings.

#### Let's avoid this



#### <u>Acknowledgements</u>

- Sexual Health Services at Waikato and Auckland DHBs
- Gay Men's Sexual Health Research Group
- New Zealand AIDS Foundation
- Hornet app

# Thank-you

joe.rich@nzaf.org.nz